



Remicade (Infliximab), Inflectra, Avsola, Renflexis Administration Orders

Patient Name: _____ DOB: _____ Weight _____ lb/kg
ICD 10 Code _____ (required)
Allergies _____

Pre-medicate 30 minutes before the start of the infusion.

Tylenol _____ mg PO Benadryl _____ mg PO/IV Other _____

Medication: _____

Induction dose of _____ mg/kg week 0,2,6 then

Maintenance dose of _____ mg/kg OR _____ mg every _____ weeks x 1 year.

- Dilute in 250ml NS for final concentration between 0.4mg/ml and 4mg/ml.
- Do not infuse other medications into IV line with Infliximab.
- Must be used within 3 hours of admixture.
- Start infusion at 20 ml/hr. for 15 minutes.
- Increase rate to 60ml/hr. for 15 minutes.
- Increase to 150ml/hr. for 60 minutes.
- Increase to 275 ml/hr. until infusion is completed.

Monitoring

- Assess patient's vitals before and after infusion.
- Monitor and record presence/absence of infusion related reactions; chills, pruritis or urticaria, chest pain. Hypotension, hypertension, or dyspnea OR hypersensitivity reaction.

PRN for nausea: Zofran 4- 8 mg IVP. If hypersensitivity or infusion related reaction occurs during infusion:

- Stop infliximab and start normal saline.
- Give Benadryl 50 mg IVP.
- Give Solu-Medrol 125 mg IVP.
- Call physician before restarting infusion.
- For anaphylaxis, give Epinephrin 1:1000, 0.5mg SQ, repeat twice at 20-minute intervals and call 911. O2 by nasal canula @2-5 L/ per minute PRN dyspnea or chest pain.

Physician Signature: _____ Date: _____

Physician Name (printed): _____

Office Number: _____

4362 Auburn Blvd., Sacramento, CA 95841
1215 Plumas St. Ste 1300B Yuba City, Ca 95991
Ph: 916-678-5840 Fax: 916-678-5839